

PATENT COOPERATION TREATY

RECEIVED
 23 JAN 2007

From the RECEIVING OFFICE

PCT

COMMUNICATION IN CASES FOR WHICH
NO OTHER FORM IS APPLICABLE

To: DR. S. BANNAN GEC HEALTH CARE AMERSHAM PLACE LITTLE CHALFONT BUCKINGHAMSHIRE HP7 9NA	Date of mailing <small>(day/month/year)</small> <div style="text-align: right; font-size: 1.2em;">22-01-07.</div>
Applicant's or agent's file reference <div style="text-align: center; font-size: 1.2em;">PA 984A</div>	REPLY DUE <div style="text-align: right;">See paragraph 1 below</div>
International application No. <div style="text-align: center; font-size: 1.2em;">PCT/GB 1999/004410</div>	International filing date <small>(day/month/year)</small> <div style="text-align: right; font-size: 1.2em;">23-12-99.</div>
Applicant <div style="text-align: center; font-size: 1.2em;">NYCOMED AMERSHAM PLC (et al)</div>	

1. ☐ REPLY DUE within _____ months/days from the above date of mailing
- ☒ NO REPLY DUE, however, see below _____
- ☐ IMPORTANT COMMUNICATION
- ☒ INFORMATION ONLY

DUE DATE	
DATE OF MAILING	✓ KE ✓ PD
DATE OF RECEIPT	SB
CASE NO.	P29848-PCT

2. COMMUNICATION:

Further to our conversation and your facsimile of the 18th January 2007 concerning the above-mentioned Pct application.

As requested, I enclose a copy of page 1 of the faxed description for your information/retention.

Finally, I confirm said page was page 1 of the faxed application ^{specification} received at the P/GB on the 23rd December 1999. (23-12-99.), this being the International filing date of said case.

Name and mailing address of Receiving Office The Patent Office Cardiff Road, Newport South Wales NP10 8QQ	Authorized officer J. R. LLOYD-THOMAS ROOM: GY19 EXT: 4340 Telephone No. 4036
---	---

IN VITRO NMR METHOD

This invention is concerned with nuclear magnetic resonance spectroscopy. The technique involves observing the spectrum of a NMR active nuclear species, particularly a hyperpolarised nucleus, in order to obtain information about the environment in which the species is present. The spectra of NMR active nuclei vary depending on their environment, as reported in the literature (PNAS, 93, 12932-6, 1996).

Noble gases having non-zero nuclear spin can be hyperpolarised, i.e. have their polarisation enhanced over the equilibrium polarisation, e.g. by the use of circularly polarised light. Preferred techniques for hyperpolarisation include spin exchange with an optically pumped alkali metal vapour and metastability exchange. Noble gases to which this technique can be applied include ^3He and ^{129}Xe . As described by M S Albert *et al* in US Patent 5,545,396, the technique can be used to prepare hyperpolarised noble gases which can then be administered by inhalation for magnetic resonance imaging of the human body.

It is known that the hyperpolarisation of a noble gas can be transferred to another NMR active species by physical contact. Thus WO 97/37239 (Lawrence Berkeley National Laboratory) describes a method which involves: contacting a sample containing an NMR active nucleus with a hyperpolarised noble gas; scanning the sample using nuclear magnetic resonance spectroscopy, magnetic resonance imaging, or both, in order to detect the NMR active nucleus. WO 98/30918 (Nycomed Imaging AS) relates to *ex-vivo* dynamic nuclear polarisation (DNP) of the NMR active nuclei of an MR imaging agent by a hyperpolarised gas where the gas is separated from the MR imaging agent prior to administration to the body.

The present invention concerns the hyperpolarisation of one or more NMR active nuclei of compounds involved in an assay. The hyperpolarisation may be carried out using a variety of techniques, such as polarisation transfer from a noble gas, "Brute force", DNP (WO 98/58272, Nycomed Imaging AS) and the para hydrogen ($p\text{-H}_2$) method, as explained below.

The transfer of hyperpolarisation according to the present invention may be achieved by using a hyperpolarised noble gas, preferably ^3He or ^{129}Xe , or a mixture of such gases, to effect